
This article was downloaded by:[USDA National Agricultural Library]

On: 16 November 2007

Access Details: [subscription number 777121083]

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



# Food Additives & Contaminants

Publication details, including instructions for authors and subscription information: http://www.informaworld.com/smpp/title~content=t713599661

# Gene profiling for studying the mechanism of aflatoxin biosynthesis in **Aspergillus flavus** and **A. parasiticus**

Jiujiang Yu <sup>a</sup>; Catherine M. Ronning <sup>b</sup>; Jeffery R. Wilkinson <sup>c</sup>; Bruce C. Campbell <sup>d</sup>; Gary A. Payne <sup>e</sup>; Deepak Bhatnagar <sup>a</sup>; Thomas E. Cleveland <sup>a</sup>; William C. Nierman <sup>bf</sup>

<sup>a</sup> USDA/ARS, Southern Regional Research Center, New Orleans, LA70124, USA

b The Institute for Genomic Research, Rockville, MD 20850, USA

<sup>c</sup> Mississippi State University, Mississippi State, MS 39762, USA

d USDA/ARS, Western Regional Research Center, Albany, CA 94710, USA

<sup>e</sup> Department of Plant Pathology, North Carolina State University, Raleigh, NC 27695, USA

<sup>f</sup> Department of Biochemistry and Molecular Biology, The George Washington University School of Medicine, Washington, DC 20037, USA

Online Publication Date: 01 October 2007

To cite this Article: Yu, Jiujiang, Ronning, Catherine M., Wilkinson, Jeffery R., Campbell, Bruce C., Payne, Gary A., Bhatnagar, Deepak, Cleveland, Thomas E. and Nierman, William C. (2007) 'Gene profiling for studying the mechanism of aflatoxin biosynthesis in **Aspergillus flavus** and **A. parasiticus**', Food Additives & Contaminants, 24:10, 1035 - 1042 To link to this article: DOI: 10.1080/02652030701513800

URL: http://dx.doi.org/10.1080/02652030701513800

#### PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: http://www.informaworld.com/terms-and-conditions-of-access.pdf

This article maybe used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.



# Gene profiling for studying the mechanism of aflatoxin biosynthesis in Aspergillus flavus and A. parasiticus

JIUJIANG YU<sup>1†</sup>, CATHERINE M. RONNING<sup>2†</sup>, JEFFERY R. WILKINSON<sup>3†</sup>, BRUCE C. CAMPBELL<sup>4</sup>, GARY A. PAYNE<sup>5</sup>, DEEPAK BHATNAGAR<sup>1</sup>, THOMAS E. CLEVELAND<sup>1</sup>, & WILLIAM C. NIERMAN<sup>2,6</sup>

<sup>1</sup>USDA/ARS, Southern Regional Research Center, 1100 Robert E. Lee Blvd., New Orleans, LA70124, USA, <sup>2</sup>The Institute for Genomic Research, 9712 Medical Center Drive, Rockville, MD 20850, USA, <sup>3</sup>Mississippi State University, Mississippi State, MS 39762, USA, <sup>4</sup>USDA/ARS, Western Regional Research Center, 800 Buchanan Street, Albany, CA 94710, USA, <sup>5</sup>Department of Plant Pathology, North Carolina State University, Raleigh, NC 27695, USA, and <sup>6</sup>Department of Biochemistry and Molecular Biology, The George Washington University School of Medicine, 2300 Eye Street NW, Washington, DC 20037, USA

(Received 12 April 2007; accepted 8 June 2007)

#### **Abstract**

Aflatoxins are toxic and carcinogenic polyketide metabolites produced by certain fungal species, including *Aspergillus flavus* and *A. parasiticus*. Many internal and external factors, such as nutrition and environment affect aflatoxin biosynthesis; therefore, we analyzed the transcriptome of *A. flavus* using expressed sequence tags (ESTs) from a normalized cDNA expression library constructed from mycelia harvested under several conditions. A total of 7218 unique ESTs were identified from 26110 sequenced cDNA clones. Functional classifications were assigned to these ESTs and genes, potentially involved in the aflatoxin contamination process, were identified. Based on this EST sequence information, a genomic DNA amplicon microarray was constructed at The Institute for Genomic Research (TIGR). To identify potential regulatory networks controlling aflatoxin contamination in food and feeds, gene expression profiles in aflatoxin-supportive media versus non-aflatoxin-supportive media were evaluated in *A. flavus* and *A. parasiticus*. Genes consistently expressed in several aflatoxin-supportive media are reported.

**Keywords:** Aspergillus flavus, mycotoxin, aflatoxin, expressed sequence tags (EST), microarray, gene profiling, aflatoxin biosynthesis, genetic regulation, pathogenesis

# Introduction

Aspergillus flavus and A. parasiticus are the two major species that produce aflatoxins (Cotty et al. 1994; Cotty 1997; Bennett and Klich 2003), the toxic and most carcinogenic natural secondary metabolites known. A. flavus is the predominant species responsible for aflatoxin contamination of crops prior to harvest or during storage (Cotty et al. 1994; Bennett and Klich 2003). The acute toxicity and carcinogenic properties of aflatoxins have been recognized for over 40 years (Lancaster et al. 1961). Due to the significant health and economic impacts of aflatoxin

contamination, the chemistry, enzymology and genetics of the aflatoxin biosynthetic pathway in *A. flavus* and *A. parasiticus* have been actively studied (Cleveland and Bhatnagar 1990; Bhatnagar et al. 1991; Bennett et al. 1997; Yabe and Nakajima 2004; Yu 2004; Yu et al. 2004b, 2005) to decipher the biological mechanism of toxin production. Studies on aflatoxin biosynthesis and its genetic regulation in *A. flavus* and *A. parasiticus* led to the cloning of 25 clustered genes within a 75-kb DNA region responsible for enzymatic conversions in the aflatoxin biosynthetic pathway (Yu et al. 2004a, 2004b). Regulatory genes within the gene cluster, such as

Correspondence: Dr. Jiujiang Yu, USDA/ARS, Southern Regional Research Center, 1100 Robert E. Lee Blvd., New Orleans, LA70124, USA. E-mail: jiuyu@srrc.ars.usda.gov

†Yu, Ronning and Wilkinson contributed equally to this work

ISSN 0265–203X print/ISSN 1464–5122 online © 2007 Taylor & Francis

DOI: 10.1080/02652030701513800

1036 J. Yu et al.

aflR and aflS (aflf), have also been cloned (Chang et al. 1993; Payne et al. 1993; Meyers et al. 1998) and their mode of action on activating aflatoxin pathway gene expression reported (Chang et al. 1995; Chang 2003). Fungal nutritional and environmental factors (Payne and Brown 1998; Yu et al. 2003; OBrian et al. 2007), as well as stage of fungal development (Calvo et al. 2002; Calvo et al. 2004), were also found to affect aflatoxin formation. However, the global regulatory control of aflatoxin formation beyond its biosynthetic gene cluster and the process of signal transduction in turning aflatoxin production on or off remains unknown. The A. flavus EST project identified 7218 unique ESTs (Yu et al. 2004c). Genes that are potentially involved in aflatoxin formation have been categorized (Yu et al. 2004c) and include those in the aflatoxin biosynthetic pathway, signal transduction/ global regulation, virulence/pathogenicity of the fungus and stress response/antioxidation (Yu et al. 2004c). We report here additional categories of genes that are potentially involved in secondary metabolism, transport or drug resistance, and fungal development or sporulation. These genes may also be important in aflatoxin formation. Microarray technology is a powerful tool in profiling genes or components that are differentially regulatory expressed under specific conditions. Previous works using microarray for profiling aflatoxin biosynthetic genes have been very successful (OBrian et al. 2003; Price et al. 2005; Wilkinson et al. 2007; OBrain et al. 2007). Price et al. (2005) using targeted 763 geneelement microarray studied the gene expression under conducive and non-conducive conditions of carbon source, nitrogen source, pH and temperature in A. parasiticus. In a more focused investigation on carbon source effects, we studied the gene expression in both A. flavus and A. parasiticus, as well as A. oryzae, under two sets of carbon source media. YES vs. YE and GMS vs. PMS. In this report, we present, in addition to the results of EST data (unpublished due to page limitation in a previous publication), the genes consistently expressed in seven experiments in several medium shift experiments in A. flavus and A. parasiticus, as well as A. oryzae.

#### Materials and methods

Fungal strains and culture conditions

A. flavus NRRL 3357 (ATCC# 20026; A. flavus SRRC 167), the wild type strain widely used in laboratory and field studies, was selected for making the EST library. To make the library as representative as possible for gene expression, fungal mycelia were grown in eight separate media and collected at five time points (Table I), as reported earlier (Yu et al. 2004c). The mycelia were harvested by filtration through miracloth at five different time points (18, 24, 36, 48, 72, 96 h) following inoculation with a conidial suspension. The harvested mycelial samples were mixed and frozen in liquid nitrogen for RNA purification.

For microarray experiments, A. flavus NRRL 3357, A. flavus #13, A. parasiticus SRRC 143 (ATCC # 56775 or SU-1) and A. oryzae RIB 40 were chosen. A. flavus NRRL 3357 is the same strain that was used for the EST (Yu et al. 2004c) and for the whole genome sequencing project (Payne et al. 2006). A. flavus #13 belongs to S strain that produces small sclerotia and large amount of personal aflatoxins (Cotty, communication). A. parasiticus SRRC 143 is a wild-type strain used widely in laboratories for molecular genetic studies on aflatoxin biosynthesis. A. oryzae RIB 40 is a food grade organism that is widely used in fermentation industry (Machida et al. 2005).

Normalized cDNA library construction, sequencing and annotation

A normalized cDNA expression library was constructed by Incyte Genomics Inc. (Palo Alto, CA, USA) with the mixed mycelia described in Table I as starting material. Single pass, unidirectional (5' end) sequencing was completed at TIGR on ABI 3700 sequencing machines. The sequence assembly was performed using the CAP3 program (Huang and Madan 1999) and Paracel Transcript Assembler (version 2.6.2, http://www.paracel.com) with modifications by the TIGR bioinformatics team (Yu et al. 2004c).

Table I. Fungal mycelia collected from cultures and at various time-points (in g).

Culture media	18 h	24 h	48 h	72 h	96 h	Aflatoxin production
Wheat bran solid	20	20	20	20	20	Supportive
GMS liquid (Glucose minimal salt)	10	10	10	10	10	Supportive
PMS liquid (Peptone minimal salt)	10	10	10	10	10	Non-supportive
PDB (Potato dextrose broth)	10	10	10	10	10	Supportive
Rice solid	_	_	_	_	10	Supportive
YES liquid	_	_	10	10	_	Supportive
GMS + oil	_	_	10	_	_	Supportive
PMS + oil	_	_	10	-	_	Supportive

Culture conditions and RNA isolation for microarray experiments

Erlenmeyer flasks containing 100 ml of the appropriate media were inoculated with 100 ul of 10<sup>8</sup> spores to obtain a final concentration of 10<sup>5</sup> spores ml<sup>-1</sup>. The cultures were maintained at 30°C with constant shaking at 150 rpm for 48 or 96 h. The four types of culture media used were veast extract (YE), yeast extract sucrose (YES), peptone minimal salt (PMS) and glucose minimal salt (GMS). The fungal mycelia were harvested by filtration through miracloth and frozen in liquid nitrogen, then ground to a fine powder in liquid nitrogen using mortar and pestle. Total RNA was extracted from approximately 100 mg of ground mycelium in 1 ml of TRIZOL® (Molecular Research Center, Inc., Cincinnati, OH, USA) according to the manufacturer's instructions. Total RNA was quantified using a ND-1000 spectrophotometer (NanoDrop Technologies, Wilmington, DE, USA) and visualized by gel electrophoresis to ensure quality.

### Microarray design and experiments

The genomic DNA amplicon microarray used in this study was constructed at The Institute for Genome Research (TIGR, Rockville, MD, USA) using sequence-specific primers designed according to A. flavus EST annotation (Yu et al. 2004c). A total of 5031 genes were arrayed at least three times, each for a total of 18078 features or spots. A gene can be represented on the array by more than one sequence, thus, features are not a direct representation of genes. The microarrays were printed at TIGR. PCR amplicons were purified using Millipore 96-well size exclusion vacuum filter plates. The purified PCR products were resuspended in water and diluted 1:1 with DMSO before printing. Using an Intelligent Automation Systems spotting robot, the PCR products were arranged in triplicate at high density Telechem Superamine aminosilane-coated microscope slides. The outline of the microarray experiments are shown in Table II. The microarray hybridization was performed by aflatoxin-producing medium vs. non-producing medium. Probe labeling by dye-flips was made to balance dye bias during labeling. No 96-h time-point *A. flavus* and *A. oryzae* samples were used in array hybridization due to degradation of RNA samples at the later growth stage. Data from the 48-h time-point from *A. parasiticus* SRRC 143 was obtained. Consistent comparative results were demonstrated only by the 48-h in *A. flavus* (*A. oryzae*) with 96-h *A. parasiticus* under different medium conditions

#### cDNA synthesis, labeling and hybridization

Three similar protocols were used in cDNA synthesis, labeling and hybridization for comparison, including The Institute for Genomic Research Aminoallyl RNA labeling protocol and microarray hybridization protocol (http://pfgrc.tigr.org/ protocols.shtml); 3DNA Array 900 protocol (Genisphere Inc., Hatfield, PA, USA); and Array 900 MPX<sup>TM</sup> protocol (Genisphere Inc., Hatfield, PA, USA). Pre-hybridization was performed according to TIGR's protocols. When using the 3DNA Array 900<sup>TM</sup> and Array 900 MPX<sup>TM</sup> kit, the post-hybridization manufacturer's instructions were followed. Each experiment consisted of one sample grown in an aflatoxin-productive condition coupled to a sample from a non-productive condition. Each hybridization was repeated with duplicate dve-flip.

### Data acquisition and analysis

Labeled and washed slides were scanned using either a ScanArray5000XL (GSI Lumonics, Packard Biochip, Packard BioScience Company, Billerica, MA, USA) or a GenePix 4000B (Axon Instruments) and the independent TIFF images from each channel were analyzed using TIGR Spotfinder (http://pfgrc.tigr.org/tools.shtml). To remove nonspecific background signals, the raw data were normalized using a local regression technique, **LOWESS** (LOcally WEighted Scatterplot Smoothing) with the MIDAS software tool (http:// pfgrc.tigr.org/tools.shtml). The LOWESS normalized dye-flips were then combined to minimize any artifacts introduced by the Cy3 or Cy5 dyes;

Table II. Microarray experimental design.

Experiment	Strain	Hybridization protocol	Time-point (h)	Array pair
1	A. flavus NRRL 3357	TIGR	48	GMS vs. PMS
2	A. oryzae RIB40	TIGR	48	GMS vs. PMS
3	A. flavus #13	Array 900	48	YES vs. YE
4	A. parasiticus SRRC 143	Array 900	96	YES vs. YE
5	A. parasiticus SRRC 143	Array 900	96	GMS vs. PMS
6	A. parasiticus SRRC 143	Array 900 MPX	96	YES vs. YE
7	A. parasiticus SRRC 143	Array 900 MPX	96	GMS vs. PMS

1038 J. Yu et al.

the resulting data were averaged over duplicate gene features on each array for each replicate experiment. Some genes are represented by two separate nonoverlapping fragments, therefore, since a gene may be detected by each fragment, they are defined as features so that gene count will not be over reported. All calculated gene expression ratios were log<sub>2</sub>transformed and aflatoxin biosynthetic gene expressions were examined by cross comparison between experiments using TIGR MeV (Saeed et al. 2003) (http://pfgrc.tigr.org/tools.shtml). To determine genes that were significantly differentially expressed across all experiments, only those genes with log<sub>2</sub>transformed expression levels equal to or higher than 1.96 standard deviations from the mean (95% confidence) were selected. This filtration of significantly expressed genes was conducted using MIDAS and the resulting lists of the genes were examined further by cross comparison between experiments using TIGR MeV (Saeed et al. 2003) (http:// www.tm4.org/mev.html).

#### Results and discussion

# EST sequences obtained

Of more than 26000 sequenced A. flavus cDNA clones, 19618 high-quality EST sequences were obtained and submitted to the NCBI GenBank database (http://www.ncbi.nlm.nih.gov/). The functional classification of these ESTs has been presented in Gene Index by The Institute for Genomic Research (TIGR, http://www.tigr.org; currently is managed by The Dana Farber Cancer Institute at the web site http://compbio.dfci.harvard.edu/tgi). The high-quality sequences were assembled into 7218 unique ESTs consisting of 3749 tentative consensus sequences (TCs) and 3469 singletons (Yu et al. 2004c). These unique ESTs represent putative expressed genes and account for an

estimated 60% of the predicted  $12\,000$  functional genes in the *A. flavus* genome (Payne et al. 2006).

# Genes coregulated with aflatoxin biosynthesis

Several categories of genes that are presumably involved in aflatoxin biosynthesis, either directly or indirectly, have been published previously (Yu et al. 2004c), such as genes associated with the aflatoxin biosynthetic pathway, global regulation and signal transduction, virulence/pathogenicity, or stress response and antioxidation. The additional categories of ESTs, having homology to sequences in the existing GenBank databases and not published in a previous paper due to page limitation, are presented in Tables III–V below.

Fungal developmental processes, such as sporulation and sclerotia formation, are related to secondary metabolite formation (Calvo et al. 2002; Chang et al. 2002), and mutants deficient in sporulation are unable to produce aflatoxins (Calvo et al. 2002). The genes involved in fungal development and conidiation were identified in the A. flavus EST library (Table III). A. flavus is known to be resistant to almost all of the antibiotics available in market, making it difficult for targeted mutagenesis using an antibiotic-resistant marker system. This feature also presents a significant obstacle for developing effective fungicides to protect crops, animals and even human beings from fungal infections. The aflatoxin pathway cluster gene aflT may be involved in aflatoxin efflux by pumping toxins out of fungal cells, has been cloned in both A. flavus and A. parasiticus (Yu et al. 2004a). However, our study indicates that its function may be insignificant to aflatoxin production or secretion (Chang et al. 2004). Many ESTs with homologies to genes that confer drug resistance have been identified (Table IV). The identification of these transporters could help explain why A. flavus is resistant to most

Table III. Genes involved in fungal development/sporulation.

EST ID	Hit accession #	Putative function	Organism	% ID	% Sim	<i>E</i> -value
NAGAZ57TV	PIR S29903	Cell wall proline-rich protein	Hedera helix	38.89	52.78	0.00012
NAGBG48TV	GB   BAA35140.1	Chitinase	Emericella nidulans	68.55	84.68	1.30E-44
TC10379	GB   AAO50865.1	Similar to Leishmania major. Ppg3	Dictyostelium discoideum	19.67	41.84	0.045
NAGBA10TV	GB   AAD34461.1	Ascus development protein 3	Neurospora crassa	36.19	54.29	2.70E-10
NAGBD49TV	SP P40552	Cell wall protein TIR3 precursor	Saccharomyces cerevisiae	27.97	45.45	4.20E-06
NAGBU14TV	SP P10169	Conidiation-specific protein 8	Neurospora crassa	41.11	51.11	5.90E-08
TC10246	SP P10169	Conidiation-specific protein 8	Neurospora crassa	41.11	51.11	5.80E-08
TC11956	SP P10713	Conidiation-specific protein 10	Neurospora crassa	71.6	80.25	2.40E-25
TC12083	SP P10713	Conidiation-specific protein 10	Neurospora crassa	67.86	73.21	7.00E-12
NAFEA74TV	GI   585141	Developmental regulator flbA	Emericella nidulans	93	96	1.00E-126
NAGBI96TV	SP O60032	Conidiophore development protein hymA	Emericella nidulans	89.66	95.4	4.50E-39
TC8878	SP O60032	Conidiophore development protein hymA	Emericella nidulans	90.83	95.83	3.30E-53

1039

Table IV. Genes involved in transport and drug resistance.

EST ID	Hit accession #	Putative function	Organism	% ID	% Sim	<i>E</i> -value
NAFCX67TV	GB   CAB46647.1	Putative multiple drug resistance protein	Penicillium olsonii	60.42	77.60	5.50E-60
NAGCB21TV	GB   CAC28822.2	Multidrug resistance-associated protein	Neurospora crassa	52.56	67.31	6.70E-30
NAGDC32TV	SP P32767	Pleiotropic drug resistance regulatory protein 6	Saccharomyces cerevisiae	36.43	57.36	1.20E-08
NAFCV74TV	GB   CAD21494.1	Conserved hypothetical protein	Neurospora crassa	48.61	70.83	4.80E-11
NAGCB47TV	GB   CAD71016.1	Probable mportin-alpha export receptor	Neurospora crassa	51.20	72.80	1.20E-27
TC11203	GB   CAD70375.1	Probable fluconazole resistance protein	Neurospora crassa	55.62	69.10	9.00E-51
TC11795	PIR   T30882	Multidrug resistance protein 1	Aspergillus flavus	100	100	0
TC9584	GB   AAA50353.1	Metal resistance protein	Saccharomyces cerevisiae	49.70	72.26	4.30E-94
TC9769	SP P32386	ATP-dependent bile acid permease	Saccharomyces cerevisiae	56.28	74.49	9.80E-75
TC10317	GB   CAD56485.1	Fructose facilitator	Zygosaccharomyces bailii	70.18	81.58	1.00E-81
TC8832	PIR   G83503	Probable MFS transporter PA1131	Pseudomonas aeruginosa	25.62	43.84	0.043

Table V. Genes potentially involved in secondary metabolism.

EST ID	Hit accession #	Putative function	Organism	% ID	% Sim	<i>E</i> -value
NAGCA18TV	SP P32637	Glyceraldehyde 3-phosphate dehydrogenase	Podospora anserina	79.67	89.43	3/E-48
NAGCA40TV	GB BAB49569.1	NADH-dependent dyhydrogenase	Mesorhizobium loti	29.63	50.62	0.038
NAGCB08TV	GB   AAF26281.1	Monooxygenase	Aspergillus parasiticus	100	100	2.30E-26
NAGCB61TV	SP O47950	NADH-ubiquinone oxidoreductase 19.3 kDa subunit	Neurospora crassa	92.42	96.21	1.60E-66
NAGCB73TV	GB   CAC87049.1	Mitochondrial NADH : ubiquinone oxidoreductase	Bos taurus	38.79	61.21	1/E-14
NAGCO63TV	GB   AAF26274.1	NADH oxidase	Aspergillus parasiticus	87.32	92.25	1.30E-60
NAGCQ74TV	PIR T38538	Probable oxidoreductase	Schizosaccharomyces pombe	43.07	64.23	4.40E-24
NAGDC76TV	GB   AAD53257.1	Dioxygenase alpha subunit	Australian soil clone OD16	28.81	49.15	0.92
TC10309	GB BAC20334.1	Monooxygenase	Aspergillus oryzae	98.69	99.22	1.20E-213
TC10899	GB AAF26280.1	Cytochrome <i>P</i> 450 monooxygenase	Aspergillus parasiticus	98.35	100	4.60E-63
TC8634	PIR S55328	Serine-type carboxypeptidase precursor	Aspergillus phoenicis	80	85.79	1.40E-79
TC8800	SP P36060	NADH-cytochrome b5 reductase precursor	Saccharomyces cerevisiae	42.72	61.17	3.70E-49
TC10233	SP P41747	Alcohol dehydrogenase I	Aspergillus flavus	99.71	99.71	2.10E-184
NAGBA01TV	GB AAD13655.1	T-2 toxin biosynthesis protein	Fusarium sporotrichioides	45.11	63.59	6.50E-32
TC8750	PIR   AF3007	Epoxide hydrolase	Agrobacterium tumefaciens	33.33	46.53	3.80E-07

of the antibiotics available and the difficulty in developing antifungal drugs effective in the filamentous fungi, especially A. fumigatus (Moore et al. 2000). Preliminary assembly of the A. flavus whole genome sequence revealed at least 65 genes encoding for ABC-2 type transporter family proteins (Payne et al. 2006); these proteins could play important roles in multi-drug resistance.

Aspergillus flavus produces many types of secondary metabolites, of which aflatoxins are the most notorious. In addition to the genes that are directly involved in aflatoxin biosynthesis, we have identified a category of genes that are putatively involved in the biosynthesis of secondary metabolites, such as those encoding polyketide synthases, fatty acid synthases,

carboxylases, dehydrogenases, reductases, oxidases, oxidoreductases, epoxide hydrolases, mono- or di-oxigenases, cytochrome P450 monooxygenases methyltransferases (Yu et al. 2004b). In addition to aflatoxin pathway genes, we have identified many ESTs that could be involved in the biosynthesis of additional secondary metabolites (Table V). Preliminary annotation of the A. flavus genome indicated that there are a number of gene clusters or genes that potentially encode for enzymes involved in secondary metabolite production (Payne et al. 2006), including 35 putative polyketide synthases, 24 putative non-ribosomal peptide synthases and over 130 cytochrome P450 enzymes (Payne et al. 2006).

Table VI. Genes expressed in common under several aflatoxin-producing carbon source media in A. flavus and A. parasiticus as detected by microarray experiments.

	1 2 3 4 5 6	1	73	3	4	ſΩ	9	7
		TIGR	TIGR	Array 900	Array 900	Array 900	Array 900MPX	Array 900MPX
		Af3357	Ao RIB $40$	Af #13	Ap 143	Ap 143	Ap 143	Ap 143
Experiment		48h	48h	48h	96 h	96 h	96 h	96 h
TIGR EST#	Putative function	GMS vs. PMS	GMS vs. PMS	YES vs. YE	YES vs. YE	GMS vs. PMS	YES vs. YE	GMS vs. PMS
NAFAY22TV	Indolepyruvate decarboxylase	3.8079698	5.1009502	3.6314514	5.8113933	3.7582371	4.3477554	2.0790656
NAFC095TV	Peptide transporter ptr2-a	-3.3024704	-4.6637373	-5.3831525	-5.291969	-5.321002	-2.8490677	-3.7928588
NAFDC06TV	Indolepyruvate decarboxylase	3.7094684	5.2063975	3.7822225	5.8895564	3.9719558	3.5608275	3.0449924
NAFDH31TV	Antigenic cell wall protein MP1	-2.991371	-4.1788764	-3.7679496	-2.872515	-4.855476	-2.4336514	-3.0006616
NAGAC91TV	RNA export mediator GLE1	1.9787863	2.296317	3.9615142	3.5454264	2.7678428	2.688219	1.4570416
NAGAG12TV	Unknown	4.1175394	4.962635	3.6760864	4.8225274	4.4131126	3.859379	2.7965271
NAGAS67TV	Unknown	4.1640916	5.1858807	3.8899755	4.9367576	4.2946796	3.6684997	3.2086217
NAGAT37TV	Aspartate aminotransferase	-4.972739	-5.4448504	-4.886115	-3.8674343	-4.5452228	-2.6529703	-2.184085

Notes: The numbers are  $\log 2$  ratios showing gene expression up-regulation (positive values) and down-regulation (negative values). Log 2 = 2 represents a 4-fold increase in gene expression;  $\log 2 = 4$  represents a 16-fold increase in gene expression, and so on.

Genes profiled through microarray analysis under aflatoxin-supportive carbon source media

Gene profiling experiments using microarrays were performed under aflatoxin-supportive in contrast to non-supportive conditions. A total of 2340 scorable expressed features were detected across all experiments. All of the aflatoxin pathway genes and associated sugar cluster genes were detected in this comparison in most, but not all experiments (data not shown). However, the aflatoxin pathway regulatory genes aflR and aflS (aflf) (Chang et al. 1993; Payne et al. 1993; Meyers et al. 1998) were not detected in any of the experiments, possibly as levels of their gene transcripts are not high enough to be detected by microarray. However, the expression levels of the regulatory genes aflR and aflS were high enough to be detected by RT-PCR and real-time PCR methods in previous experiments performed in this laboratory (data not shown). Even under non-aflatoxin-supportive medium condition (such as PMS), a very low level of aflR transcript was detected by the sensitive RT-PCR procedure in A. flavus (unpublished data and Chang, personal communication). The aflR transcript level and, hence, the AfIR regulator protein under the nonsupportive condition is too low to activate the transcription of aflatoxin biosynthetic pathway structural genes for producing aflatoxins. After filtration of expressed genes at the 95% confidence level and examination by cross comparison using TIGR MeV program, five genes that were consistently up-regulated in response to the carbon source and three genes down-regulated, including two genes with unknown function, were identified as being in common for all of the culture conditions (Table VI). It is apparent that these genes are most likely involved in primary metabolism rather than directly involved in secondary metabolism. These genes could be coregulated and co-expressed with aflatoxin pathway genes under aflatoxin-producing conditions. It could also represent shift from one metabolic pathway to another, more favourable for secondary metabolite formation, such as aflatoxins. More importantly, these genes could be the links between primary and secondary metabolism. The exact nature of biological involvements of these up-and down-regulated genes can only be deciphered by gene knock-out experiments. It is also interesting to find that these genes are consistently expressed in A. oryzae RIB 40, the non-aflatoxin producer used in food fermentation, indicating that the three species, A. flavus, A. parasiticus and A. oryzae share a common biological pathway response to carbon-rich medium. The nontoxigenicity of *A. oryzae* is most likely due to loss of ability to produce aflatoxins at a later stage of its regulatory hierarchy.

#### Conclusion

EST and microarray analyses are robust tools for profiling genes involved in specific biological processes. However, for certain low level expressed genes such as regulatory genes, more sensitive protocols are needed for future research. The genes identified in this study by EST and microarray techniques are putative candidates for further investigation. Without additional biological evidence, it is very difficult to predict their exact biological functions. Traditional targeted mutagenesis and high throughput gene knock-out systems are necessary to elucidate the functions of a large number of genes in the post-genome era.

#### Acknowledgements

This paper was approved for publication as Journal Article No. J-11066 of the Mississippi Agricultural and Forestry Experiment Station, Mississippi State University.

#### References

Bennett JW, Chang PK, Bhatnagar D. 1997. One gene to whole pathway: The role of norsolorinic acid in aflatoxin research. Advances in Applied Microbiology 45:1–15.

Bennett JW, Klich M. 2003. Mycotoxins. Clinical Microbiology Reviews 16:497–516.

Bhatnagar D, Cleveland TE, Kingston DG. 1991. Enzymological evidence for separate pathways for aflatoxin B1 and B2 biosynthesis. Biochemistry 30:4343–4350.

Calvo AM, Wilson RA, Bok JW, Keller NP. 2002. Relationship between secondary metabolism and fungal development. Microbiology and Molecular Biology Reviews 66:447–459.

Calvo AM, Bok J-W, Brooks W, Keller NP. 2004. VeA is required for toxin and sclerotial production in *Aspergillus parasiticus*. Applied and Environmental Microbiology 70:4733–4739.

Chang PK. 2003. The Aspergillus parasiticus protein AFLJ interacts with the aflatoxin pathway-specific regulator AFLR. Molecular Genetics and Genomics 268:711–719.

Chang PK, Cary JW, Bhatnagar D, Cleveland TE, Bennett JW, Linz JE, Woloshuk CP, Payne GA. 1993. Cloning of the *Aspergillus parasiticus apa-2* gene associated with the regulation of aflatoxin biosynthesis. Applied and Environmental Microbiology 59:3273–3279.

Chang PK, Ehrlich KC, Yu J, Bhatnagar D, Cleveland TE. 1995. Increased expression of *Aspergillus parasiticus aflR*, encoding a sequence-specific DNA-binding protein, relieves nitrate inhibition of aflatoxin biosynthesis. Applied and Environmental Microbiology 61:2372–2377.

Chang PK, Bennett JW, Cotty PJ. 2002. Association of aflatoxin biosynthesis and sclerotial development in *Aspergillus parasiticus*. Mycopathologia 153:41–48.

1042 J. Yu et al.

- Chang PK, Yu J, Yu JH. 2004. *aflT*, a MFS transporter-encoding gene located in the aflatoxin gene cluster, does not have a significant role in aflatoxin secretion. Fungal Genetics and Biology 41:911–920.
- Cleveland TE, Bhatnagar D. 1990. Evidence for de novo synthesis of an aflatoxin pathway methyltransferase near the cessation of active growth and the onset of aflatoxin biosynthesis in *Aspergillus parasiticus* mycelia. Canadian Journal of Microbiology 36:1–5.
- Cotty PJ. 1997. Aflatoxin-producing potential of communities of *Aspergillus* section *Flavi* from cotton producing areas in the United States. Mycological Research 101:698–704.
- Cotty PJ, Bayman DS, Egel DS, Elias KS. 1994. Agriculture, aflatoxins and *Aspergillus*. In: Powell K, editor. The Genus *Aspergillus*. New York: Plenum Press. pp 1–27.
- Huang X, Madan A. 1999. CAP3: A DNA sequence assembly program. Genome Research 9:868–877.
- Lancaster MD, Jenkins FP, Philip JM. 1961. Toxicity associated with certain samples of groundnuts. Nature 192:1095–1096.
- Machida M, Asai K, Sano M, Tanaka T, Kumagai T, Terai G, et al. 2005. Genome sequencing and analysis of Aspergillus oryzae. Nature 438:1157–1161.
- Meyers DM, OBrian G, Du WL, Bhatnagar D, Payne GA. 1998. Characterization of *afl*<sup>3</sup>, a gene required for conversion of pathway intermediates to aflatoxin. Applied and Environmental Microbiology 64:3713–3717.
- Moore CB, Sayers N, Mosquera J, Slaven J, Denning DW. 2000.
  Antifungal drug resistance in Aspergillus. Journal of Infection 41:203–220
- OBrian GR, Fakhoury AM, Payne GA. 2003. Identification of genes differentially expressed during aflatoxin biosynthesis in *Aspergillus flavus* and *Aspergillus parasiticus*. Fungal Genetics and Biology 39:118–127.
- OBrian GR, Georgianna DR, Wilkinson JR, Yu J, Abbas HK, Bhatnagar D, Cleveland TE, Nierman W, Payne GA. 2007. The effect of elevated temperature on gene transcription and aflatoxin biosynthesis. Mycologia 99:232–239.
- Payne GA, Brown MP. 1998. Genetics and physiology of aflatoxin biosynthesis. Annual Review of Phytopathology 36:329–362.
- Payne GA, Nystrom GJ, Bhatnagar D, Cleveland TE, Woloshuk CP. 1993. Cloning of the afl-2 gene involved in aflatoxin biosynthesis from Aspergillus flavus. Applied and Environmental Microbiology 59:156–162.

- Payne GA, Nierman WC, Wortman JR, Pritchard BL, Brown D, Dean RA, Bhatnagar D, Cleveland TE, Machida M, Yu J. 2006. Whole genome comparison of Aspergillus flavus and A. oryzae. Medical Mycology 44 Suppl:9–11.
- Price MS, Conners SB, Tachdjian S, Kelly RM, Payne GA. 2005. Aflatoxin conducive and non-conducive growth conditions reveal new gene associations with aflatoxin production. Fungal Genetics and Biology 42:506–518.
- Saeed AI, Sharov V, White J, Li J, Liang W, Bhagabati N, et al. 2003. TM4: A free, open-source system for microarray data management and analysis. Biotechniques 34:374–378.
- Wilkinson JR, Yu J, Bland JM, Nierman WC, Bhatnagar D, Cleveland TE. 2007. Amino acid supplementation reveals differential regulation of aflatoxin biosynthesis in *Aspergillus flavus* NRRL 3357 and *Aspergillus parasiticus* SRRC 143. Applied and Environmental Microbiology 74:1308–1319.
- Yabe K, Nakajima H. 2004. Enzyme reactions and genes in aflatoxin biosynthesis. Applied Microbiology and Biotechnology 64:745–755.
- Yu J. 2004. Genetics and biochemistry of mycotoxin synthesis. In: Arora DK, editor. Fungal biotechnology in agricultural, food, and environmental applications. New York: Marcel Dekker. pp 343–361.
- Yu J, Mohawed SM, Bhatnagar D, Cleveland TE. 2003. Substrate-induced lipase gene expression and aflatoxin production in *Aspergillus parasiticus* and *Aspergillus flavus*. Journal of Applied Microbiology 95:1334–1342.
- Yu J, Bhatnagar D, Cleveland TE. 2004a. Completed sequence of aflatoxin pathway gene cluster in *Aspergillus parasiticus*. FEBS Letters 564:126–130.
- Yu J, Chang PK, Ehrlich KC, Cary JW, Bhatnagar D, Cleveland TE, Payne GA, Linz JE, Woloshuk CP, Bennett JW. 2004b. Clustered pathway genes in aflatoxin biosynthesis. Applied and Environmental Microbiology 70:1253–1262.
- Yu J, Whitelaw CA, Nierman WC, Bhatnagar D, Cleveland TE. 2004c. Aspergillus flavus expressed sequence tags for identification of genes with putative roles in aflatoxin contamination of crops. FEMS Microbiology Letters 237:333–340.
- Yu J, Cleveland TE, Nierman WC, Bennett JW. 2005. Aspergillus flavus genomics: Gateway to human and animal health, food safety, and crop resistance to diseases. Revista Iberoamericana Micologia 22:194–202.